



<http://www.entm.purdue.edu/Entomology/ext/targets/newslett.htm>

May 7, 2004 - No. 8

## In This Issue

### Insects, Mites, and Nematodes

- Black Cutworm Scouting Should Begin
- Armyworm in Wheat
- Black Cutworm Adult Pheromone Trap Report
- Black Light Trap Catch Report

### Weeds

- Poison Hemlock - The Toxic Parsnip

### Plant Diseases

- Update on Risk of *Fusarium* Head Blight of Southern Indiana

### Agronomy Tips

- Fearmonger Alert: Freeze Injury Potential for Early-Planted Corn
- Requirements for Uniform Germination and Emergence of Corn
- The Emergence Process in Corn
- Corkscrewed Corn Seedlings
- Germination Events in Corn
- Wheat Condition and the Potential of Cold Injury to the Wheat Plant

### Weather Update

- Temperature Accumulations

## Insects, Mites, and Nematodes

**Black Cutworm Scouting Should Begin** – (John Obermeyer and Larry Bledsoe)

- Southern Indiana should be looking for leaf feeding and initial cutting.
- Scouting guidelines and rescue insecticides given below.
- Rotary hoe may improve control where soil soils are dry.

A few reports of cutworm damage have been received from southern Indiana, most of those are likely dingy cutworm. The dingy cutworm overwinters as a partially grown larva, so early leaf feeding is not uncommon. Fortunately it is not the threat to stand losses that the black cutworm is. Black cutworm moth catches in pheromone traps have indicated several intensive flights into the state over the last several weeks. We have begun to track heat unit accumulations from those significant flights to predict black cutworm development and beginning of plant cutting, see "Weather Update" in this issue of the *Pest&Crop*. Monitoring for black cutworm damage should begin in southern counties.

Scout by inspecting 20 consecutive plants in each of 5 areas of a field (100 plants) for cutworms and feeding activity. Count and record the number of plants cut or damaged and determine the percentage of plants affected. Also collect black cutworm larvae and determine the average instar stage. While sampling, also record how many leaves are fully unrolled (the collar of the leaf is visible on a fully unrolled leaf). Control of black cutworm may be necessary if 3 to 5% of the plants are damaged and the average larval instar is from 4 to 6. Use the following management guidelines and instar guide. Suggested foliar insecticides for control of economic infestations are listed below.

To increase the probability that adequate control will be achieved when dry soil conditions are noted, a rotary hoe may prove useful. This should stir up the soil and increase the likelihood that the cutworms will come in contact with the insecticide. Additionally, the use of a higher rate of the insecticide in 20 gallons or more water per acre may help the level of control. On no-till fields, where hoeing is not possible, applying the insecticide in the early evening may increase control, as the worms more toward the soil surface during the nighttime hours.

### Black Cutworm Management Guidelines

Average Instar 6 or of BCW	Number of Plant Leaves Fully Emerged					
	more	5	4	3	2	1
4.0	1% +	2% +	2% +	2% +	3% +	4% +
5.0	2% +	3% +	4% +	4% +	6% +	25% +
6.0	4% +	7% +	9% +	17% +	Don't	Don't
7.0	6% +	15% +	50% +	Don't	Don't	Don't

1. Look down the column at the left labeled "Average Instar of BCW" until you find the average instar of BCW found in the field. This column is called the Instar Row.
2. Look across the top of the table and find the number that best represents the "Number of Plant Leaves Fully Emerged" for the plants inspected. A leaf is fully emerged if the leaf collar is visible. The column of figures below this is called the Leaf Column.
3. Follow the Instar Row and the Leaf Column to the place where they intersect. This figure is the control threshold. If the percentage of cut or damaged plants in the field equals or exceeds this number, treatment may be advisable.

### Black Cutworm Instar Guide

Instar	Head Capsule	How to use the instar guide: Immobilize the larva by holding it with a forceps, by placing it in alcohol, or by grasping it tightly behind the head. Hold the larva flat against the paper and move it down until the head just fits inside one of the "keystone" figures. That is the most probable instar for that larva.
3	■	
4	■	
5	▲	
6	▲	
7	▲	

### Insecticides Suggested for Foliar Application to Control Cutworms in Corn

Material	Amount Per Acre and Formulation*
bifenthrin (Capture) <sup>1,2</sup>	2.1 – 6.4 fl. oz. EC
chlorpyrifos (Lorsban) <sup>1,2</sup>	1 – 2 pt. 4E
cyfluthrin (Baythroid 2) <sup>1,2</sup>	0.8 – 1.6 fl. oz. E
esfenvalerate (Asana XL) <sup>1,2</sup>	5.8 – 9.6 fl. oz. EC
lambda-cyhalothrin (Warrior) <sup>1,2</sup>	1.92 – 3.20 fl. oz. CS
methyl parathion (Penncap-M) <sup>1,2</sup>	4 pt. FM
permethrin (Ambush) <sup>1,2</sup> (Pounce) <sup>1,2</sup>	6.4 – 12.8 fl. oz. EC 4 – 8 fl. oz. 3.2EC
zeta-cypermethrin (Mustang Max) <sup>1,2</sup>	1.3 – 2.8 fl. oz. EW

\* Under dry conditions, where rotary hoeing may be needed to increase cutworm control, use the higher insecticide rate.

<sup>1</sup> Restricted Use Insecticide

<sup>2</sup> Bee Caution



Different sizes of black cutworm larvae



Seedling that has been cut below the ground

• • P&C • •

**Armyworm in Wheat** - (John Obermeyer and Larry Bledsoe)

- Larvae seen in Indiana wheat fields.
- Wheat defoliation and head clipping can result.

Jon Neufelder, Posey County CES, informs us that some armyworm damage is occurring in wheat. Armyworm moths have been making their way into the state for the last several weeks and females have been laying eggs on grassy-type plants. Hatching larvae then proceed to feed on grassy plants and crops. Fortunately the moth numbers captured in black light traps are nothing compared to the “memorable” 2001 season.

Pest managers should examine wheat plants in different areas of a field, especially where plant growth is dense. Look for flag leaf feeding, clipped heads, and armyworm droppings (excrement) on the ground. Shake the plants and count the number of armyworm on the ground and under plant debris. On sunny days, the armyworm will take shelter under crop residue or soil clods. If counts average approximately 5 or more per linear foot of row, the worms are less than 1-1/4 inches long and not parasitized or diseased, and leaf feeding is evident, control may be justified. If a significant number of armyworm are present and they are destroying the leaves, or the heads, treat immediately.

• • P&C • •

Black Cutworm Adult Pheromone Trap Report							
Week 1 = 4/22/04 - 4/28/04    Week 2 = 4/29/04 - 5/5/04							
County	Cooperator	BCW Trapped		County	Cooperator	BCW Trapped	
		Wk 1	Wk 2			Wk 1	Wk 2
Adams	Roe/Price Ag Services	8	8	Marshall	Shanks/Plymouth Pioneer (2)	7	6
Allen	Gynn/South Wind Farm	1	4	Marshall	Shanks/Plymouth Pioneer (3)	7	6
Benton	Babcock/Jasper Co. Co-op	3	3	Newton	Babcock/Jasper Co. Co-op	3	4
Clay	Smith/Growers Co-op (Brazil)	0	0	Putnam	Nicholson Consulting	29*	19*
Clay	Smith/Growers Co-op (Clay City)	1	16*	Randolph	Boyer/Davis-Purdue Ag Center	5	8
Elkhart	Kauffman/Crop Tech Inc.	1	7	Randolph	Derek Calhoun	0	18*
Fayette	Schelle/Spring Valley Farms	9	1	Rush	Tacheny/Pioneer Hi-Bred	21*	24*
Fountain	Hutson/Purdue CES	0	0	Shelby	Gabbard/Shelby Co. CES	9*	4
Fountain	Mroczkiewica/Syngenta	5	14	Sullivan	Smith/Growers Co-op (New Lebanon)	4	12*
Gibson	Hirsch Farms	2	6	Sullivan	Smith/Growers Co-op (Sullivan E)	12*	3
Greene	Maruszewski/Worthington Pioneer	9	1	Sullivan	Smith/Growers Co-op (Sullivan W)	11*	5
Johnson	Kessler/Ag Excel	-	-	Tippecanoe	Obermeyer/Purdue CES	47*	16
Knox	Smith/Growers Co-op (Fritchton)	0	0	Tipton	Johnson/Pioneer	0	2
Knox	Smith/Growers Co-op (Oaktown)	0	0	Vermillion	Hutson/Purdue CES	0	0
Lake	Kliene Farms (1)	5	5	Vigo	Smith/Growers Co-op (Terre Haute)	0	4
Lake	Kliene Farms (2)	10*	7	Warren	Babcock/Jasper Co. Co-op	2	4
Marshall	Barry/Fulton-Marshall Co-op	0	20*	White	Reynolds/Vogel Popcorn	10	1
Marshall	Shanks/Plymouth Pioneer (1)	2	0	Whitley	Walker/NE-Purdue Ag Center	2	10*

\* = Intensive Capture... an intensive capture occurs when 9 or more moths are caught over a 2-night period.

Black Light Trap Catch Report - (John Obermeyer)														
County/Cooperator	4/20/04 - 4/26/04							4/27/04 - 5/3/04						
	VC	BCW	ECB	SWCB	CEW	FAW	AW	VC	BCW	ECB	SWCB	CEW	FAW	AW
Dubois/SIPAC		2					7							
Jennings/SEPAC							2		1					4
Knox/SWPAC														3
LaPorte/Pinney Ag Center							14							11
Lawrence/Feldun Ag Center		1					24		1					12
Randolph/Davis Ag Center							1							2
Tippecanoe/TPAC Ag Center														
Vermillion/Hutson														1
Whitley/NEPAC							149							

VC = Variegated Cutworm, BCW = Black Cutworm, ECB = European Corn Borer, SWCB = Southwestern Corn Borer, CEW = Corn Earworm, FAW = Fall Armyworm, AW = Armyworm

# Weeds

**Poison Hemlock – The Toxic Parsnip** – (Glenn Nice, Bill Johnson, Tom Bauman, and Thomas Jordan)

We often get questions about wild carrot (*Daucus carota* L.) only to find out that the question is actually about poison hemlock (*Conium maculatum* L.). Although these two plants may look similar, poison hemlock is toxic to cattle, horses, swine, sheep, goats, dogs, and people when ingested. The plant produces volatile alkaloids coniine (an alkaloid similar in effect to nicotine) and gamma-coniine. The easiest way to tell the two plants apart is that poison hemlock will have purple spots or blotches on its smooth (hairless) ridged stems. Wild carrot will usually have a covering of hairs.

**Description:** Poison hemlock can often be found along roadsides, edges of cultivated fields, railroad tracks, stream banks, waste areas, and sometimes along the fence rows of pastures. Like wild carrot, poison hemlock is a biennial. This means that it lives its life over two years. In the first year, poison hemlock goes through vegetative growth. In the second year, it will produce small white flowers arranged in umbrella-like cluster, similar to wild carrot. It is in the second year, when it bolts and flowers, that it tends to catch the eye. The flower stalks can grow 3 to 8 feet tall. The leaves are finely divided having a triangular shape. When comparing both poison hemlock and wild carrot leaves, wild carrot has a more rounded lobe in the leaf. Where as Poison hemlock's leaf reminds me of a sharp arrow head. For more information and pictures of poison hemlock's description go to <[www.ppws.vt.edu/scott/weed\\_id/coima.htm](http://www.ppws.vt.edu/scott/weed_id/coima.htm)> and <[www.vet.purdue.edu/depts/add1/toxic/plant28.htm](http://www.vet.purdue.edu/depts/add1/toxic/plant28.htm)>.

**Symptoms of Poisoning:** All parts of the plant can be toxic. Young leaves in the spring are the most toxic and the root the least toxic. The fruit is most dangerous in the fall. Lethal doses can be small, so it is important not to let animals graze or feed on poison hemlock. In the case of horses, 4 to 5 pounds of the leaves may be lethal. One to 2 pounds can be lethal for cattle and 4 to 8 oz for sheep. Young animals are more susceptible. Symptoms may appear within 1 hour of ingestion. This starts with a nervous stimulation and can progress in 2 to 3 hours later into respiratory paralysis. In rare cases the animal may have convulsions. In many cases symptoms include, bloating, incoordination, intestinal irritation, dilation of pupils, rapid and weak pulse, loss of appetite, salivation, and blue coloration about the mouth. Ingestion of poison hemlock in days 55 to 75 of gestation may result in birth defects.

**Treatment:** If an animal becomes poisoned by poison hemlock, a veterinarian may administer nerve and heart stimulants as soon as possible. Large doses of

mineral oil and purgatives are also prescribed to empty the digestive tract. For more information about poison hemlock's toxic characteristics please see <[www.vet.purdue.edu/depts/add1/toxic/plant28.htm](http://www.vet.purdue.edu/depts/add1/toxic/plant28.htm)>.

**Control:** Control of poison hemlock in grass pastures is more effective in the first year of its life cycle. In a grass pasture, 2,4-D, Banvel/Clarity (dicamba), or Crossbow (2,4-D and triclopyr) provide control. Crossbow is slightly better than both 2,4-D and Banvel/Clarity. However, be aware that these herbicides will damage any legumes. Spot treatments of glyphosate products (Glyphomax Plus, Roundup WeatherMax, Touchdown, etc.) will also control poison hemlock, but be aware that this will also damage any desired vegetation. Always read and follow labels when using herbicides.

We have also noted that poison hemlock is beginning to invade no-till corn and soybean fields. Herbicides that have activity on this weed and that can be used before planting soybeans are 2,4-D, dicamba, and glyphosate. Dicamba and glyphosate have shown slightly better efficacy than 2,4-D. The best overall control would likely be attained with a mixture of glyphosate and dicamba.

**Considerations Using Dicamba in Soybean:** Use 1/2 pt./A on coarse soils and 1 pt./A on medium or fine soils with at least 2% organic mater. Not all dicamba products have PRE-plant labels in soybean: consult the label before buying for this purpose. Clarity can be applied 14 days before planting if 8 fl. oz./A or less is used and at least 1 inch of rainfall or over head irrigation occurs; however, if 16 fl. oz./A is used there is a 28 days waiting period before planting soybean.

**Considerations Using 2,4-D in Soybean:** Use 1 to 2 pt./A of a LVE (Low Volatile Ester formulation). Before planting soybean, using 2,4-D at 1 pt./A requires a 7 day waiting period before planting and if more than 1 pt./A is used, a 30 day waiting period must be observed. These restrictions may be slightly different depending on the product, please read the specific products label before buying for this purpose.

**Considerations Using Glyphosate:** Use 0.75 lb. ae/A.<sup>1</sup>

For more information on toxic plants of Indiana to livestock and pets see <[www.vet.purdue.edu/depts/add1/toxic/cover1.htm](http://www.vet.purdue.edu/depts/add1/toxic/cover1.htm)>. For online pesticide labels go to <[www.cdms.net](http://www.cdms.net)> or <[www.greenbook.net](http://www.greenbook.net)>

<sup>1</sup> ae stands for acid equivalent. This is the glyphosate weight in its acid form. Most labels will give ae per gallon to calculate amount to be used.

# Plant Diseases

## Update on Risk of *Fusarium* Head Blight of Southern Indiana - (Gregory Shaner)

The predicted risk of *Fusarium* head blight of wheat, assuming a flowering date of 3 May 2004 depends on whether there is corn residue in a wheat field. Where there is no residue, the risk is low throughout Indiana. Where corn residue is present, the model shows moderate

risk in a roughly rectangular area bordered on the west by Greene Co., on the east by Bartholomew Co., on the south by Washington Co., and on the north by Johnson Co. It is unlikely that any wheat in this area is flowering yet. Some wheat in far southwest Indiana may be flowering, but in that area the model indicates low risk. The model can be viewed at <[www.wheatcab.psu.edu](http://www.wheatcab.psu.edu)>.

# Agronomy Tips

## Fearmonger Alert: Freeze Injury Potential for Early-Planted Corn - (Bob Nielsen)

Corn planting has been proceeding at a record pace in Indiana thus far in the 2004 growing season. Reasonably warm soil temperatures throughout April have also encouraged faster emergence than usually occurs with such early-planted corn. Such early planting and emergence of corn is always at higher calendar risk of injury by frost events or lethal cold temperatures.

Of these two risk factors, lethal cold temperature is the more worrisome one since a corn plant's growing point region is relatively protected from the effects of simple frost while it remains below the soil surface. Lethal cold temperatures (28°F or less) can penetrate the upper inch or two of soil, especially dry surface soils, and kill plant tissue directly, including coleoptiles and growing points. Non-lethal injury by cold temperatures may cause deformed elongation of the mesocotyl or physical damage to the coleoptile in non-emerged seedlings, resulting in the proverbial "cork-screw" symptom and subsequent leafing out underground.

Air temperatures in northern areas of Indiana dipped to the low 30's early in the morning of 3 May, with lower-lying areas likely less than 30°F. Given the risk of frost or chilling injury to young corn; it would behoove growers to monitor early-planted fields over the next week to determine whether such injury has occurred and whether replanting may be warranted.

### Select References:

Nielsen, R.L. (Bob). 2004. Corkscrewed Corn Seedlings. Corny News Network, Purdue Univ. Available online at <[www.agry.purdue.edu/ext/corn/news/articles.04/Corkscrew-0501.html](http://www.agry.purdue.edu/ext/corn/news/articles.04/Corkscrew-0501.html)>. (Verified 5/2/04).

## Requirements for Uniform Germination and Emergence of Corn - (Bob Nielsen)

- Successful germination & emergence require adequate moisture, temperature and seed-to-soil contact.

Rapid, uniform germination and emergence of corn help set the stage for maximum grain yield at the end of the season. Without such a successful start to the season, the crop is behind the proverbial "eight-ball" right from the beginning. The good news is that there are only three simple requirements for uniform germination and emergence of corn. The bad news is that one or more of the requirements are sometimes absent from one field to another.

*Adequate and uniform soil moisture at the seed zone.* Adequate soil moisture is most simply defined as not too dry and not too wet. Most growers know what "adequate" looks and feels like. Uneven soil moisture in the seed zone can be caused by variable soil characteristics, tillage patterns, unusual weather conditions and uneven seeding depth. Uneven soil moisture in the seed zone is the primary cause of uneven emergence, the results of which can easily be yield losses of 8 to 10 percent.

**Useful Tip:** When seedbed conditions are dry, make sure that your choice of seeding depth ensures uniformly adequate soil moisture for the germination of the seed. Even though a 1.5 to 2 inch seeding depth is a good choice for many conditions, don't hesitate to increase seeding depth to 2.5 to 3 inches if that is where the uniform soil moisture is located. Planting shallower than 1.5 inches increases the risk of poor or uneven germination during subsequent drying of surface soils.

*Adequate and uniform soil temperature at the seed zone.* Adequate soil temperature is most simply defined

as being greater than 50°F at the 2-inch depth. Corn will not germinate or emerge quickly or uniformly when soil temperatures are less than 50°F. When soils warm to the mid-50°F or greater, emergence will occur in seven days or less if soil moisture is adequate.

Uneven soil temperature can be caused by soil characteristics, uneven residue cover in reduced tillage systems and uneven seeding depth control. Temperature variability is most critical when average soil temperatures are barely within the desired minimum 48 to 50°F range for corn germination.

**Useful Tip:** Dark-colored soils will typically warm more quickly than light-colored soils. If soils dry differently across the field, the drier areas will typically warm faster than the wet areas. Uneven residue cover in reduced tillage systems causes significantly lower soil temperatures under the heavier cover than under barer spots in the field. Uneven seeding depth exposes deeper planted seeds to slightly cooler seed zones than seeds placed shallower.

**Adequate and uniform seed-to-soil contact.** In order for the kernel to absorb moisture quickly and uniformly, soil must be firmed completely around the kernel. Seed-to-trash contact results from “hair-pinning” of surface trash into the seed furrow during no-till planting when soil and / or trash are too wet for adequate coulter cutting action. Seed-to-clod contact results from planting into cloddy fields created by working soil too wet. Seed-to-rock contact is, needless to say, not good for proper germination either. Seed-to-air contact results from open planter furrows when no-till planting into excessively wet soils. Germination of kernels lying in open planter furrows is dependent on rainfall keeping the open furrow environment moist.

**Useful Tip:** Whippers, wipers, movers, fingers, and other similar trash management gadgets for the planter are most beneficial when you are challenged with rocky, cloddy, or trashy surface soil conditions. They help clear the way (literally) for the planter’s double-disc openers to more easily do their job of creating an optimum seed furrow. Other planter attachments that help press the kernels into the seed furrow can improve seed-to-soil contact and seeding depth uniformity when seedbed conditions are otherwise challenging.

• • P&C • •

### The Emergence Process in Corn - (Bob Nielsen)

• Understanding the process helps you troubleshoot problems with emergence.

Successful germination alone does not guarantee successful emergence of the crop. The coleoptile must reach the soil surface before its internal leaves emerge

from the protective tissue of the coleoptile. Growth stage VE refers to emergence of the coleoptile or first leaves through the soil surface.

As with all of corn growth and development, germination and emergence are dependent on temperature, especially soil temperature. Corn typically requires from 100 to 150 GDD (growing degree days) to emerge. Under warm soil conditions, the calendar time from planting to emergence can be as little as 5 to 7 days. Under cold soil conditions, it can easily take up to four weeks to emerge.

Elongation of the **mesocotyl** elevates the coleoptile towards the soil surface. The mesocotyl is the tubular, white, stemlike tissue connecting the seed and the base of the coleoptile. Technically, the mesocotyl is the first internode of the stem.




**Useful Tip:** Physiologically, mesocotyls have the capability to lengthen from at least a 6-inch planting depth. Realistically, corn can be planted at least three inches deep if necessary to reach adequate moisture.

As the coleoptile nears the soil surface, exposure of the mesocotyl to the red light portion of the solar radiation



spectrum halts mesocotyl elongation. Continued expansion of the leaves inside the coleoptile ruptures the coleoptile tip, allowing the first true leaf to emerge above the soil surface. Since the depth at which the mesocotyl senses red light is fairly constant, the resulting depth of the crown (base) of the coleoptile is nearly the same (1/2 to 3/4 inch) at seeding depths of one-inch or greater.

**Useful Tip:** When corn is seeded very shallow (less than about 1/2 inch), the crown of the coleoptile will naturally be closer to the soil surface if not right at the surface. Subsequent development of the nodal root system can be restricted by exposure to high temperatures and dry surface soils.



Closeup of corn coleoptile


### Troubleshooting Considerations

Several factors can cause the coleoptile to split prematurely, allowing the leaves to emerge underground. Usually, more than one of the following factors are present when this problem occurs, making it difficult to place the blame on any one factor.

**Exposure to light** at deeper soil depths than usual due to cloddy seedbeds, dry seedbeds, sandy soils, or open slots in no-till.

**Injury from certain herbicides**, particularly under stressful environmental conditions. Symptoms include corkscrewed coleoptile, swollen mesocotyl and true leaves emerged from side of coleoptile.

**Surface crusting, cloddy seedbeds, rocky seedbeds, planter furrow compaction, or otherwise dense surface soil** that physically restricts mesocotyl elongation and coleoptile penetration. The pressure of the expanding



Inner leaves 'teased' out of coleoptile



Emergence of inner leaves through split tip of coleoptile



Successful emergence



Split in coleoptile side due to expansion of inner leaves



View of kernel, mesocotyl, coleoptile and leaf emergence just below the soil surface



Coleoptile tissue removed to reveal inner leaves of plumule

leaves within the coleoptile eventually ruptures the side of the coleoptile. Symptoms include corkscrewed coleoptile, swollen mesocotyl and true leaves emerged from side of coleoptile. Note the similarity to those symptoms from herbicide injury.

**Cold temperature injury**, either from exposure to long periods of soil temperatures around 50°F or from exposure to wide daily swings (25 to 30°F) in soil temperatures. Symptoms include absence of emerged coleoptile, corkscrewed mesocotyl or coleoptile and true leaves emerged from side of coleoptile. Note the similarity to those symptoms from herbicide injury.

**Useful Tip:** The mesocotyl should remain firm, white and healthy through at least the 6-leaf stage, if not longer. If it is mushy, discolored, or damaged prior to this stage, then it is likely part of the crop problem being investigated.

• • P&C • •

**Corkscrewed Corn Seedlings** - (Bob Nielsen)

- Deformed, corkscrewed, curved development of a corn plant's mesocotyl or coleoptile can be caused by several factors.

Emergence of corn occurs by the elongation of the mesocotyl that elevates the coleoptile (or "spike") to the soil surface. The mesocotyl is that white stem-like plant part located between the kernel and the crown of the coleoptile.

Mesocotyl elongation of early-planted corn occasionally veers from its usual upwardly mobile path and instead corkscrews below ground. The end result of such spiraling sub-surface seedlings is either underground leaf emergence or eventual death of the seedling. The good news is that the extent of the problem is usually limited to a few fields each year and a small number of plants (several thousand or less per acre) within an affected field.



As is usual with crop problems, several possible causes of corkscrewed seedlings exist. The challenge is to identify which is the most likely cause for any given situation.

**Restricted Emergence:** Corkscrewed mesocotyl/coleoptile development can result when the coleoptile encounters resistance as the mesocotyl elongates. Severe soil crusting or otherwise dense soil surface and cloddy soil surfaces can cause such resistance. A combination of severe sidewall compaction plus press wheel compaction over the furrow can also restrict coleoptile emergence.

**Kernel Position in Furrow:** The position of the kernel in the furrow with respect to the embryo face directly influences the initial location where the coleoptile emerges. The coleoptile, the protective covering for the plumule leaves, emerges from the embryo side of the kernel and moves toward the dent end of the kernel by virtue of the elongation of the mesocotyl. If the kernel lands with the embryo face down in the furrow, the coleoptile emerges on the bottom side of the kernel, elongates horizontally until the mesocotyl 'clears' the end of the kernel, then finally begins its upward ascent. Such an "upside-down" beginning might contribute to a seedling's susceptibility to other corkscrewing causal factors.




**Herbicide Injury:** Certain herbicides, notably cell growth inhibitors, can affect seedling shoot development especially if weather or soil conditions are not conducive for rapid growth. Quite often when herbicide is part of the blame, cool soils and significant soil crusting are also contributing factors.

**Temperature Response:** Some years ago, I came across an article from Rhodesia (Buckle & Grant. 1974. Rhod. J. Agric. Res. 12: 149-161) that described the same phenomenon and attributed it to large fluctuations between day and night soil temperatures. In their research, abnormal mesocotyl and/or coleoptile development occurred most frequently when soil temperatures fluctuated from daytime highs of about


80°F to nighttime lows of about 55°F. The data also suggested that extended periods of cold temperatures stunted and distorted seedling growth.

**Fig's C and D: Abnormal growth with wide fluctuations in soil temperature.**



Reports of corkscrewed seedlings in Indiana are usually few and far between, but occasionally correspond to situations when unusually cool soil temperatures or a dramatic fluctuation in soil temperatures occur in fields prior to seedling emergence. Some might ask how often do such dramatic fluctuations occur during the germination/emergence period? I don't have an answer for that, but can document that such a fluctuation occurred in the past several weeks. Over a six day period, beginning April 11, soil temperatures at the Purdue Agronomy Farm ranged from a daily high of 67°F to a daily low of 38°F three days later to a daily high of 79°F three days after that.

Daily Min and Max Soil Temperatures, April 2004  
WestCentral Indiana, 4-Inch Depth



Oh, by the way, I've also received my first report of corkscrewed seedlings from an early April planted field of corn in central Indiana...

• • P&C • •

## Germination Events in Corn - (Bob Nielsen)

- Understanding the process helps you troubleshoot problems with germination.

Germination is the renewal of enzymatic activity that results in cell division and elongation and, ultimately, embryo emergence through the seed coat. Germination is triggered by absorption of water through the seed coat. Corn kernels must absorb (imbibe) about 30% of their weight in water before germination begins. Less than optimum absorption of water (perhaps due to a rapidly drying seed zone) may slow or stop germination. Repeated wetting/drying cycles can decrease seed viability.

By comparison, soybeans must imbibe about 50% of their weight in water. But since soybeans are approximately 2/3 the weight of corn kernels, the total amount of absorbed water required for germination is relatively similar.

The visual indicators of germination occur in a distinct sequence. The **radicle root** emerges first, near the tip end of the kernel, within two to three days in warm soils with adequate moisture. In cooler or drier soils, the radicle root may not emerge until one to two weeks after planting.

The **coleoptile** (commonly called the 'spike') emerges next from the embryo side of the kernel within one to many days of the appearance of the radicle, depending on soil temperature. The coleoptile initially negotiates its way toward the dent end of the kernel by virtue of the elongation of the mesocotyl. The coleoptile is a rigid piece of plant tissue that completely encloses the four to five embryonic leaves (plumule) that formed during grain development of the seed production year. The plumule leaves slowly enlarge and eventually cause the coleoptile to split open as it nears the soil surface.

The **lateral seminal roots** emerge last, near the dent end of the kernel.

### Troubleshooting Considerations


When temperatures are optimum, these three parts of the seedling may emerge from the kernel on nearly the same day. Excessively cool soils may delay the appearance of the coleoptile and lateral seminal roots for more than a week after the radicle root emerges. It is not uncommon in cold planting seasons to dig seed two weeks after planting and find only short radicle roots and no visible coleoptiles.



Appearance of radicle root 3 days after planting (DAP)




Appearance of radicle root 3 DAP (different plant)



Appearance of plumule and lateral seminal roots 4 DAP



Swelling of plumule area 4 DAP (different plant)



Appearance of plumule, lateral seminal roots and radicle root 4 DAP (different plant)

When excessively cold and/or wet soils delay germination and/or emergence, the kernel and young seedling are subjected to lengthier exposure to damaging factors such as soil-borne seed diseases, insect feeding and injury from pre-plant or pre-emergent herbicides and carryover herbicides from a previous crop

**Select References:**

Hardman, L.L. and J.L. Gunsolus. 1998. Corn Growth and Development & Management Information for Replant Decisions. Univ. of Minnesota Ext. Service Pub. No. FO-05700. Available online at <[www.extension.umn.edu/distribution/cropsystems/DC5700.html](http://www.extension.umn.edu/distribution/cropsystems/DC5700.html)>. (Verified 5/1/04).  
 Ritchie, S.W., J.J. Hanway, and G.O. Benson. How a Corn Plant Grows. Iowa State Univ. Extension Sp. Rpt. 48. Available online at <<http://maize.agron.iastate.edu/corngrows.html>>. (Verified 5/1/04).


Don't forget, this and other timely information about corn can be viewed at the Chat 'n Chew Café on the Web at <[www.kingcorn.org/cafe](http://www.kingcorn.org/cafe)>. For other information about corn, take a look at the Corn Growers' Guidebook on the Web at <[www.kingcorn.org](http://www.kingcorn.org)>.

• • P&C • •


**Wheat Condition and the Potential of Cold Injury to the Wheat Plant** – (Ellsworth P. Christmas and Charles W. Mansfield)

- Wheat condition very good across the entire state.
- Low temperatures and potential damage to wheat.

The Indiana Agricultural Statistics Service in their 'Indiana Crop and Weather Report' for the week ending on May 2 rated 86% of the wheat crop as good to excellent. They report that 80% of the wheat crop is jointed and 11% is headed.



Same as Image 4 except with seed coat removed to reveal plumule and lateral seminal roots



Appearance of plumule, lateral seminal roots and radicle root 4 DAP (different plant)

On the morning of May 3 and/or 4, 7 stations across northern and central Indiana reported low temperatures between 26 and 31°F. Six stations reported temperature of 26 or 31°F on May 3 and five on May 4, 2004.

The stage of growth of the wheat varied from head stage in southwestern Indiana to early joint stage in northern Indiana. Once wheat has reached the boot stage, temperatures at or below 28°F for a period of two or more hours can result in freeze damage to the wheat plant. Symptoms of this injury may include floret sterility, head entrapment, stem damage or leaf discoloration. From emergence of the head until early milk stage, temperatures at or below 30°F for a period of two or more hours can result in freeze damage to the wheat plant. This damage can be characterized by floret sterility, white awns or heads, stem damage or leaf

discoloration. The extent of the injury is dependent on the actual temperature and length of time the plant was subjected to the low temperature.

The temperatures reported above may not represent the actual temperatures in the field at the level of the developing wheat heads. It is not unusual for the temperature at the level of the heads to be 4 to 5 degrees colder than the temperatures at the official height of the recording thermometer.

It is not possible to confirm freeze damage to wheat until the plant has had at least 5 days to continue to grow and develop. Table 1 below gives a summary of the temperature required to cause freeze injury to the wheat plant at various stage of growth and the symptom most likely to be evident.

<b>Growth stage</b>	<b>Approx. injurious temperature (two hours)</b>	<b>Primary symptoms</b>	<b>Yield effect</b>
Tillering	12F (-11C)	Leaf chlorosis; burning of leaf tips; silage odor; blue cast to field	Slight to moderate
Jointing	24F (-4C)	Death of growing point; leaf yellowing or burning; lesions, splitting, or bending of lower stems; odor	Moderate to severe
Boot	28F (-2C)	Floret sterility; head trapped in boot; damage to lower stem; leaf discoloration; odor	Moderate to severe
Heading	30F (-1C)	Floret sterility; white awns or wheat heads; damage to lower stem; leaf discoloration	Severe
Flowering	30F (-1C)	Floret sterility; white awns or white heads; damage to lower stem; leaf discoloration	Severe
Milk	28F (-2C)	White awns or white heads; damage to lower stems; leaf discoloration; shrunken, roughened, or discolored kernels	Moderate to severe
Dough	28F (-2C)	Shriveled, discolored kernels; poor germination	Slight to moderate

# Weather Update

Temperatures as of May 5, 2004

HU48 = heat units at a 48°F base from Jan. 1, for alfalfa weevil development (begin scouting at 200)

HU50 = heat units at a 50°F base from date of intensive moth capture, for black cutworm development (larval cutting begins about 300)

GDD(5) = Growing Degree Days from April 7 (5% of Indiana's corn planted), for corn growth and development

GDD(42) = Growing Degree Days from April 21 (42% of Indiana's corn planted), for corn growth and development


## 4" Bare Soil Temperatures 5/5/04

MAP KEY			
Location			
HU48	HU50	GDD(5)	GDD(42)

**Bug Scout says, "Hey northern Indiana don't forget the alfalfa crop...look for weevil damage!"**



**Bug Scout says, "Southern Counties should be scouting seedling corn for cutworm damage."**



Location	Max.	Min.
Wanatah	67	48
Winamac	64	49
Chalmers	54	51
Bluffton	52	46
W Laf Agro	64	51
Tipton	58	50
Farmland	57	47
Crawfordsville	56	52
Liberty	67	47

The **Pest&Crop** is produced by the Departments of Agronomy, Botany and Plant Pathology, and Entomology at Purdue University. The Newsletter is published monthly February, March, October, and November. Weekly publication begins the first week of April and continues through mid-September. If there are questions or problems, contact the Extension Entomology Office at (765) 494-8761. Reference to products in this publication is not intended to be an endorsement to the exclusion of others which may have similar uses. Any person using products listed in this publication assumes full responsibility for their use in accordance with current directions of the manufacturer.

**DISCLAIMER**

It is the policy of the Purdue University Cooperative Extension Service, David C. Petritz, Director, that all persons shall have equal opportunity and access to the programs and facilities without regard to race, color, sex, religion, national origin, age, marital status, parental status, sexual orientation, or disability. Purdue University is an Affirmative Action employer. This material may be available in alternative formats 1-888-EXT-INFO (398-4636)