

# Vegetable Insects

Department of Entomology

## MANAGING STRIPED CUCUMBER BEETLE POPULATIONS ON CANTALOUPE AND WATERMELON

Ricky E. Foster, Extension Entomologist

The striped cucumber beetle feeds exclusively on cucurbits (i.e., cucumber, squash, watermelon and cantaloupe). This insect is responsible for more insecticide applications on cantaloupes than any other pest in Indiana.

Both the adult beetle and the immature stages (larvae) feed on watermelon and cantaloupe. Larvae feed on roots and stems at or below ground level. This feeding is seldom noticed, but can kill seedlings and reduce the growth of larger plants. Adults feed on the leaves and stems of the plant and can be found feeding under the canopy, or at the base of the plant. Although adults feeding on the stems can kill young plants, the most severe damage is caused by the transmission of a bacterium, *Erwinia tracheiphila* (E.F. Smith) Holland, that causes bacterial wilt in cucurbits.

Striped cucumber beetles are approximately 1/3 inch long with black heads, yellowish bodies, and 3 black stripes running the length of their wing covers. Western corn rootworm adults look remarkably similar to striped cucumber adults. However, striped cucumber adults have black abdomens, while western corn rootworm have yellow abdomens.



Striped cucumber beetles (Photo Credit: Dan Egel)

### LIFE CYCLE

Beetles overwinter as adults on edges of fields or in woods under litter or other plant debris. In late April or early May, beetles begin to emerge from their overwintering areas and feed on wild cucumbers and tree blossoms. Once watermelon or cantaloupe is planted, the beetles find and rapidly infest these preferred hosts. This initial infestation is usually the largest of the season. Beetles can be found feeding on plants within 24 hours after transplanting. The length of infestation is variable, running from a few days to a few weeks, depending on weather and other factors. In northern parts of Indiana, the beetles' appearance is usually more gradual over the entire season. Therefore, beetles may be found in moderate numbers at any time during the growing season.

Early in the season, beetles feed extensively on leaves and stems. During this early part of the season, beetles mate, and females lay eggs in the soil at the base of the plant. The yellow, oval eggs hatch in 5-8 days into white, worm-like larvae that feed on the roots. Larval feeding lasts for 14-20 days and can kill, or stunt seedlings and small plants. The larvae then pupate underground and emerge as adults from late June to mid-July (late July in northern Indiana). Adults remain active until fall. The second generation adults usually have little impact on plant growth, but may feed on developing fruit, making scars or pits in the rind that render the fruit unmarketable.

### TRANSMISSION OF BACTERIAL WILT

Adults, but not larvae, carry a bacterium (*Erwinia tracheiphila*) within their gut and on their mouthparts that causes bacterial wilt in susceptible cucurbits. Cantaloupe and cucumber are very susceptible to bacterial wilt, but watermelon is not. Beetles harbor the bacterium in their gut over the winter.

The bacterium is introduced into the plant only when the beetles feed. Once the plant is "wounded," bacteria enter the wound either from the beetles' mouthparts (seldom during the early season) or its feces (most likely source of infection). Very few beetles actually are able to infect a plant, although the



Severe wilt in field (Photo Credit: Dan Egel)

percentage increases through the season, and depends on the amount of time beetles are allowed to feed on the plant (Table 1). Therefore, early detection of beetles is essential for good bacterial wilt control. Even so, when populations of beetles reach densities of 30 to 100 per plant, many plants will become infected with the bacterium in a relatively short time.

Month	% Beetles Able to Infect Plants
May	0.7 - 1.2%
June	1.1 - 1.6%
July	3.2 - 3.8%
August	7.1 - 8.5%
Time (Hr) Beetles Allowed to Feed on Plant	% of Plants Infected
0	0%
24	1.83%
48	4.53%
72	10.67%

Bacteria are not capable of actively infecting plants. They must move into the plants through wounds and by free water. Water must be present for large percentages of plants to become infected. Therefore, rain or dew at the time of intense beetle feeding can significantly increase the incidence of bacterial wilt.

Once plants are infected, there is no way to stop the progression of the disease. The bacteria multiply in the water conducting tubes of the plant and stop the flow of water. Plants begin to wilt anywhere between 7-21 days after infection.

The disease can be diagnosed in the field by cutting through the stem at the base of a wilted plant and holding the two cut ends together for 10 seconds. As they are slowly pulled apart, a white viscous “sap” will appear as a string between the two cut ends. This stringy sap is actually the bacterial slime in the water-conducting tubes.



Confirmation of infection by bacterial wilt pathogen. Top photo is a before and bottom is after pulled apart. (Photo Credit: Chris Gunter)

**MONITORING BEETLE POPULATIONS**

After plants are set in the field, they should be watched closely for the first appearance of the beetles. Beetles usually are worse along the field edge, but large groups of them can be found randomly throughout the field. Therefore, the entire field needs to be walked to find these pockets of beetles. Beetles usually come into a field rapidly, and within 8-12

hours a field with very few beetles can become a field with 40 beetles per plant. Because the incidence of bacterial wilt infection is greater when beetles feed longer, it is essential to detect the early population surge within 24 hours of the initial beetle build-up.

Once this initial surge decreases, weekly monitoring should detect any fluctuations in beetle numbers. The second generation usually does not require treatment unless beetles are damaging the fruit.

## MANAGEMENT

### Soil Insecticides

Furadan, Admire, and Platinum can be applied at planting for control of striped cucumber beetles on cucurbits. Furdana 4F is available in Indiana as a Special Local Needs registration, and the user must have a copy of the supplemental label when using this product. These three products will provide 2-4 weeks of control, depending on rainfall and rate of plant growth. However, these products should not be counted on to provide complete control when striped cucumber beetle populations are high. Foliar applications of insecticides may also be needed. All of these soil insecticides will provide some level of control of seedcorn maggots, although none provide complete control. Admire and Platinum will also provide control of aphids.

Studies have shown a large increase in early watermelon yields and, to some extent, cantaloupe yields, when Furadan

is applied as a soil insecticide. Early watermelon yields can be increased by as much as 20 - 50% when Furadan is used at planting. Cantaloupe yields may be increased by as much as 10-20%. These increases probably are the result of plant growth regulator activity by carbofuran, combined with controlling larval feeding on roots.

### Foliar Sprays on Watermelon

Because watermelon is not susceptible to bacterial wilt, insecticides should only be applied when an average of 5 or more beetles are feeding on each plant (Table 2). Once the initial early flush of beetles is gone, any further foliar insecticide applications should be unnecessary, unless they are feeding in large numbers on developing fruit. Growers should be aware that use of some foliar insecticides may increase problems with aphids and mites.

### Foliar Sprays On Cantaloupe

Cantaloupes are **very** susceptible to bacterial wilt and, therefore, little feeding damage can be tolerated. Once a plant is infected with the bacterium, the disease cannot be controlled and the plant will die. The only way to avoid the disease is to prevent beetles from feeding on the plants. Plants should be watched closely for the first wave of beetles any time after transplanting. If an average of more than 1 beetle per plant is observed feeding on cantaloupes, an insecticide should be applied. Growers can expect 3 days of control from Sevin XLR and 5-7 days from the other insecticides recommended.

**Table 2. Foliar insecticides recommended for striped cucumber beetle control on watermelon and cantaloupe**

Insecticide	Amount Per Acre	PHI (Days)	Comments
Asana XL	5.8 to 9.6 fl. oz.	3	Do not exceed 0.25 lb. per season.
Baythroid	2.4 - 2.8 fl. oz.	0	Do not exceed 11.2 fl. oz. per acre or 4 applications per season.
Capture/Brigade	2 EC at 2.6 - 6.4 fl. oz.	3	Do not apply more than 19.2 fl. oz. per acre per year.
Danitol	2.4 EC at 10.67 - 16 fl. oz.	7	Do not exceed 42.67 fl. oz. per acre per season.
Mustang Max	2.8 - 4 fl. oz.	1	Do not exceed 24. fl. oz. per acre pr season.
Pounce	3.2 EC at 4 - 8 fl. oz. or 25 WP at 6.4 - 12.8 oz.	0	Cantaloupe only, apply a minimum of 4 gal. finished spray/acre by air or 20 gal. finished spray/acre with ground equipment.
Sevin	XLR Plus at 1 qt. 80 SP at 1 1/4 lb.	3	Some phytotoxicity may result when carbaryl is applied during hot, humid weather, especially on seedlings and newly set plants. Carbaryl is highly toxic to bees visiting blooms.

READ AND FOLLOW ALL LABEL INSTRUCTIONS. THIS INCLUDES DIRECTIONS FOR USE, PRECAUTIONARY STATEMENTS (HAZARDS TO HUMANS, DOMESTIC ANIMALS, AND ENDANGERED SPECIES), ENVIRONMENTAL HAZARDS, RATES OF APPLICATION, NUMBER OF APPLICATIONS, REENTRY INTERVALS, HARVEST RESTRICTIONS, STORAGE AND DISPOSAL, AND ANY SPECIFIC WARNINGS AND/OR PRECAUTIONS FOR SAFE HANDLING OF THE PESTICIDES.

Revised 4/2008

It is the policy of the Purdue University Cooperative Extension Service that all persons have equal opportunity and access to its educational programs, services, activities, and facilities without regard to race, religion, color, sex, age, national origin or ancestry, marital status, parental status, sexual orientation, disability or status as a veteran. Purdue University is an Affirmative Action institution. This material may be available in alternative formats.

1-888-EXT-INFO (398-4636)

<<http://www.ces.purdue.edu/new>>