

# Pest&Crop newsletter

**Purdue Cooperative Extension Service and USDA-NIFA Extension IPM Grant**

This work is supported in part by Extension Implementation Grant 2017-70006-27140/ IND011460G4-1013877 from the USDA National Institute of Food and Agriculture

## In This Issue

- [Handy Bt Trait Table Update](#)
- [Purdue Crop Performance Program Trial Results](#)
- [Sulfur Fertilizer Response of Corn - Background and Research](#)

## Update

- [Average Temperature: Departure from Mean Nov. 6 - Dec. 5, 2018](#)
- [Accumulated Precipitation \(in\): Departure from Mean Nov. 6- Dec. 5, 2018](#)

It is the policy of the Purdue University that all persons have equal opportunity and access to its educational programs, services, activities, and facilities without regard to race, religion, color, sex, age, national origin or ancestry, marital status, parental status, sexual orientation, disability or status as a veteran. Purdue is an Affirmative Action Institution. This material may be available in alternative formats. 1-888-EXT-INFO Disclaimer: Reference to products in this publication is not intended to be an endorsement to the exclusion of others which may have similar uses. Any person using products listed in this publication assumes full responsibility for their use in accordance with current directions of the manufacturer.

Pest&Crop newsletter © Purdue University - [extension.entm.purdue.edu/newsletters/pestandcrop](http://extension.entm.purdue.edu/newsletters/pestandcrop)  
Editor: Tammy Luck | Department of Entomology, Purdue University, 901 W. State St., West Lafayette, IN 47907

# Handy Bt Trait Table Update

**Author: John Obermeyer**

A wonderful reference to understand which Bt-traited corn has efficacy against specific insects is the “Handy Bt Trait Table.” New in this update

are the known insects resistant to Bt toxins. This table, produced by Chris DiFonzo, Field Crops Entomologist at Michigan State University, can be downloaded [HERE](#).

---

It is the policy of the Purdue University that all persons have equal opportunity and access to its educational programs, services, activities, and facilities without regard to race, religion, color, sex, age, national origin or ancestry, marital status, parental status, sexual orientation, disability or status as a veteran. Purdue is an Affirmative Action Institution. This material may be available in alternative formats. 1-888-EXT-INFO Disclaimer: Reference to products in this publication is not intended to be an endorsement to the exclusion of others which may have similar uses. Any person using products listed in this publication assumes full responsibility for their use in accordance with current directions of the manufacturer.

Pest&Crop newsletter © Purdue University - [extension.entm.purdue.edu/newsletters/pestandcrop](http://extension.entm.purdue.edu/newsletters/pestandcrop)  
Editor: Tammy Luck | Department of Entomology, Purdue University, 901 W. State St., West Lafayette, IN 47907

# Purdue Crop Performance Program Trial Results

**Author: John E. Woodmansee, Purdue Extension Educator, Agriculture/Natural Resources**

COLUMBIA CITY, Ind. — The Purdue Crop Performance Program (PCPP) annual Corn and Soybean Performance Trial Bulletin was recently posted online. The corn and soybean results are compiled from the 2018 growing season. Phil Devillez, program director, reported that over 40 trial results were collected from 13 test locations across Indiana. The PCPP is designed to help farmers and other agriculturalist compare crop data across Indiana.



The Purdue Crop Performance Program is designed to help farmers and other agriculturalist compare crop data across Indiana and the surrounding states. (USDA)

photo by Lance Cheung via Flickr)

The Purdue Crop Performance Program is designed to help farmers and other agriculturalist compare crop data across Indiana and the surrounding states. “What makes this program unique is the multi-state corn or soybean databases,” said Devillez. “The databases are a tool used to effectively search and compare crop test results.” The databases are distinctive to Purdue University with results from The Ohio State University, Michigan State University, University of Kentucky and the University of Illinois.

For purposes of the report, the state is split into three regions, North, Central and South. In the north and central regions early, mid and late tests were established based on appropriate maturities. The tests established in the south were early and late.

Another feature of the report is a 2018 Indiana Rainfall Summary, including April through September rainfall totals in various regions, all compared to a 30-year average. For example, at the Columbia City data collection point, the April through September total in 2018 was 17.05 inches, compared with the 30-year average of 21.10 inches. Other northern Indiana data collection sites received from 20.23 to 22.21 inches in 2018 during the same time period.

In evaluating Indiana data, there are approximately 20 corn and 20 soybean companies that are partnered with the program and over 200 corn hybrids and 180 soybean varieties evaluated each year. Each year a corn and soybean bulletin is designed to describe this data. If you are interested in joining Purdue’s program or have questions contact Phil Devillez, program director.

Find the report at PCPP’s website:

<https://ag.purdue.edu/agry/PCPP/Pages/default.aspx>.

It is the policy of the Purdue University that all persons have equal opportunity and access to its educational programs, services, activities, and facilities without regard to race, religion, color, sex, age, national origin or ancestry, marital status, parental status, sexual orientation, disability or status as a veteran. Purdue is an Affirmative Action Institution. This material may be available in alternative formats. 1-888-EXT-INFO Disclaimer: Reference to products in this publication is not intended to be an endorsement to the exclusion of others which may have similar uses. Any person using products listed in this publication assumes full responsibility for their use in accordance with current directions of the manufacturer.

Pest&Crop newsletter © Purdue University - [extension.entm.purdue.edu/newsletters/pestandcrop](http://extension.entm.purdue.edu/newsletters/pestandcrop)  
Editor: Tammy Luck | Department of Entomology, Purdue University, 901 W. State St., West Lafayette, IN 47907

# Sulfur Fertilizer Response of Corn – Background and Research Update

**Authors: Jim Camberato and Bob Nielsen**

## Summary

Sulfur (S) deficiency is becoming more common in Indiana crops because S emissions from coal-fired power plants have decreased over the past few decades and, thus, so has atmospheric S deposition. We conducted 11 large plot strip trials at 7 locations to examine corn yield response to S applied as ammonium thiosulfate (ATS) in starter and/or sidedress nitrogen fertilizers. Grain yield was increased by sidedress S in 6 of 11 trials. At responsive sites, yield increases ranged from 4 to 22 bu/acre and averaged 14 bu/acre. Sulfur in starter fertilizer only, increased corn grain at only one location.

## Sulfur for crops from air and soil

In the past, atmospheric deposition of sulfur (S) from the burning of coal provided enough S to satisfy crop needs. Pollution controls and conversion of power plants from coal to natural gas have greatly reduced S deposition, increasing the possibility of S deficiency in crops grown in Indiana. Indeed, we have diagnosed S deficiency in corn, soybean, wheat, and alfalfa in both research fields and farmer's fields. With limited S from the atmosphere, the most important source of S in most situations is S released from organic matter (OM). Each percent OM in the plow layer contains about 100 lb S/acre. Organic S must be mineralized to sulfate – S ( $\text{SO}_4\text{-S}$ ) to be taken up by crop plants, in much the same way that organic nitrogen (N) is made available to crop plants. At most only a few percent of the organic S is made available to the crop annually – approximately 3 lb S/acre per percent OM or less.



Figure 1. Shelby County sulfur (S) response trial in 2018. 16-row strips of light green corn can be seen where no S was applied to the corn crop. Where S was applied, the corn had darker green color.

Since mineralization is a process carried out by living microorganisms, soil temperature and moisture largely determine when and how much of the organic S is made available to the crop. Cold and excessively wet or dry conditions reduce microbial activity and reduce S availability from soil organic matter and crop residues.

Sulfate – S is relatively mobile in most soils (similar to nitrate) because it has a double negative charge and is repelled by the negative charge of the soil, unlike nutrients such as potassium, calcium, or magnesium which are attracted to soil particles. Although  $\text{SO}_4\text{-S}$  can bind to iron and aluminum in the soil, these elements are much more likely to bind phosphate at the exclusion of  $\text{SO}_4\text{-S}$  and as a result  $\text{SO}_4\text{-S}$  is easily leached from soils, especially sandy soils.

The occurrence of S deficiency within a field may be highly variable since soil S availability varies considerably with soil OM and texture. Sulfur deficiency is often seen in sandier, lower OM, and higher elevation areas of a field while lower lying, higher OM, and heavier textured areas may have sufficient S. Soil testing methods measure the  $\text{SO}_4\text{-S}$  form of S. Unfortunately soil testing is not particularly useful for predicting S deficiency because it does not take into account the organic S component that might become available to the crop. Additionally, the  $\text{SO}_4\text{-S}$  component that is actually measured may also be leached from the soil between the time of sampling and the time of crop need.

## Fertilizer materials

There are several fertilizers available for correcting S deficiency (Table 1). Adding ammonium thiosulfate to urea – ammonium nitrate solutions or blending ammonium sulfate with urea or other granular fertilizers are convenient and cost effective ways to provide S in a timely manner to corn. Otherwise, sulfate-of-potash-magnesia (sul-po-mag or K-mag) or potassium sulfate can be blended with muriate of potash to provide S and potassium (K). The inclusion of magnesium in sul-po-mag may be an extra benefit compared to potassium sulfate if soil magnesium levels are low. Generally these fertilizers are spread prior to planting therefore some of the  $\text{SO}_4\text{-S}$  might be lost from sandy soils before the time of crop need.

Naturally-occurring mined gypsum and several by-product sources of gypsum can be applied to provide S as well. Some forms of pelletized gypsum can be blended with other fertilizers or a ground gypsum can be applied with a lime spreader. Unless pelletized, however, higher than necessary rates of S will be applied with gypsum, which is difficult to spread at rates less than 500 to 1000 pounds per acre (85 to 170 pounds of S per acre assuming 17% S). If carryover of S occurs, the S will be utilized in later years. However, in sandy soils, where leaching is likely, the benefit in future years may be minimal.

Elemental S must be oxidized by soil bacteria to  $\text{SO}_4\text{-S}$  before being plant available. Warm temperatures and good moisture and aeration are required for S-oxidizing bacteria to function. Sulfur oxidation is minimal at soil temperatures less than 50°F. Even at 75°F oxidation of S is about 15% of that at 85°F, so peak rates of S oxidation do not occur until late spring. Since the availability of elemental S may be minimal in early spring, a fertilizer containing all or at least some  $\text{SO}_4\text{-S}$ , in addition to elemental S, is preferred over a fertilizer with elemental S alone.

**Table 1. Sulfur-containing fertilizers and their approximate composition. the actual nutrient concentration of any particular fertilizer should be expressed on the guarantee.**

Fertilizer	%N	%K <sub>2</sub> O	%S	%Mg	%Ca
Ammonium sulfate	21	0	24	0	0
Ammonium thiosulfate	12	0	26	0	0
Elemental sulfur	0	0	>90	0	0
Gypsum (calcium sulfate)	0	0	19	0	23
Potassium magnesium sulfate	0	22	23	11	0
Potassium sulfate	0	50	18	0	0

## Summary of sulfur response trials

In 2017 and 2018 we conducted 11 trials exploring the impact of starter and/or sidedress S on corn yield (Figure 2). To provide S we mixed

ammonium thiosulfate (ATS) with urea ammonium nitrate (UAN-32% or 28%) or a mixture of UAN and ammonium polyphosphate (10-34-0) as a starter fertilizer in 2x2 placement or as ATS with UAN as a sidedress application. Rates of S ranged from 5 lb/acre as starter to 25 lb S/acre as sidedress.

Sulfur application increased yield in 6 of 11 trials. Yield increases were 4, 8, 13, 18, 20, and 22 bushels per acre at the responsive sites (Fig. 2). Sidedress rates of 10 or 15 pounds of S per acre at sidedress maximized the yield response. There was no additional benefit to higher rates of S. Sulfur in the starter fertilizer at a low rate (<5 lb S/acre) increased yield (4 bu/acre) at only one location. At this one location, there was no additional benefit of sidedress S. We have not conducted enough experiments yet to determine the frequency and magnitude of S deficient conditions across Indiana. Nor do we know with certainty the soil and management conditions most likely to result in S deficiency. Consider conducting simple strip trials on your farm to determine whether you need additional S to maximize yield and profit. If you are interested in working with us on these trials please do not hesitate to contact us.




Figure 2. Yield benefit to sidedress S in Indiana in 2017 (above line) or 2018 (below line). ne=no trial that year.

*Acknowledgements: Indiana Corn Marketing Council, Pioneer Hybrid Intl., Beck's Hybrids, Purdue Agricultural Centers, and farmer cooperators. This work was supported by the USDA National Institute of Food and Agriculture, Hatch Project 1010713.*


It is the policy of the Purdue University that all persons have equal opportunity and access to its educational programs, services, activities, and facilities without regard to race, religion, color, sex, age, national origin or ancestry, marital status, parental status, sexual orientation, disability or status as a veteran. Purdue is an Affirmative Action Institution. This material may be available in alternative formats. 1-888-EXT-INFO Disclaimer: Reference to products in this publication is not intended to be an endorsement to the exclusion of others which may have similar uses. Any person using products listed in this publication assumes full responsibility for their use in accordance with current directions of the manufacturer.

Pest&Crop newsletter © Purdue University - [extension.entm.purdue.edu/newsletters/pestandcrop](http://extension.entm.purdue.edu/newsletters/pestandcrop)  
 Editor: Tammy Luck | Department of Entomology, Purdue University, 901 W. State St., West Lafayette, IN 47907



# Average Temperature: Departure from Mean Nov. 6 – Dec. 5, 2018

Average Temperature (°F): Departure from Mean  
November 6, 2018 to December 5, 2018



It is the policy of the Purdue University that all persons have equal opportunity and access to its educational programs, services, activities, and facilities without regard to race, religion, color, sex, age, national origin or ancestry, marital status, parental status, sexual orientation, disability or status as a veteran. Purdue is an Affirmative Action Institution. This material may be available in alternative formats. 1-888-EXT-INFO Disclaimer: Reference to products in this publication is not intended to be an endorsement to the exclusion of others which may have similar uses. Any person using products listed in this publication assumes full responsibility for their use in accordance with current directions of the manufacturer.

Pest&Crop newsletter © Purdue University - [extension.entm.purdue.edu/newsletters/pestandcrop](http://extension.entm.purdue.edu/newsletters/pestandcrop)  
Editor: Tammy Luck | Department of Entomology, Purdue University, 901 W. State St., West Lafayette, IN 47907

# Accumulated Precipitation (in): Departure from Mean Nov. 6-Dec. 5, 2018

Accumulated Precipitation (in): Departure from Mean  
November 6, 2018 to December 5, 2018

