

Beekeeping

Department of Entomology

WHAT BEEKEEPERS SHOULD KNOW ABOUT BEE MITES

Greg J. Hunt and Krispn Given, Extension Entomology

THE TWO PARASITIC BEE MITES CURRENTLY IN THE U.S.

The tracheal mite (*Acarapis woodii*) was first described as the causal agent of Isle of Wight disease in the United Kingdom in the early 20th century. U.S. beekeepers first noticed these mites in their hives about 1990. The tracheal mites were the first to cause heavy winter losses of bee hives in the United States. However, recent surveys show that Varroa mites cause more colony losses in North America and Europe than any other factor.

The Varroa mite (*Varroa destructor*) was originally a parasite of the Asian honey bee (*Apis cerana*) but gained the ability to infest the western honey bees commonly used for beekeeping (*Apis mellifera*) and is now found worldwide.

TRACHEAL MITES

Unlike Varroa mites which are ectoparasites, tracheal mites are microscopic parasites that live inside the tracheae in the bodies of bees (Figure 1). The tracheae are the breathing tubes of insects that distribute oxygen throughout the tissues in their bodies. Female mites emerge from the thoracic trachea and infest bees that are generally less than 24 hours old. Older bees are resistant to infestation. Tracheal mites partly block the breathing passages. Their presence



Figure 1. Tracheal mites inside of the thoracic tracheae of a honey bee (400X magnification). (Photo Credit: Lilia de Guzman, USDA ARS)

and feeding can seriously affect bee health and lead to the death of the colony. The problem is worse in colder climates and during the winter when bees cluster tightly together and colony populations are reduced. However as a result of natural selection and breeding, most North American bee stocks are now sufficiently resistant to tracheal mites so that mite treatments are generally not needed to prevent colony losses. It should be noted that some fumigants that are used to control Varroa mites such as thymol and formic acid, also kill tracheal mites.

To detect tracheal mites one must slice the thorax (the part where the wings attach) of each bee into pieces, stain the tissue, and examine it under a microscope at 400 times magnification. If less than 5% of the bees are infested, the hive should be able to survive the winter. Beekeepers rarely check for tracheal mites or use treatments to control them, because resistant bees are now common.

VARROA MITES

Biology

Varroa mites look like ticks on honey bees (Figure 2). The mites appear as coppery brown (or red-brown) discs that are about 1.5 mm wide and they are quite mobile. The mites pierce the bee's exterior to feed. Varroa mites reproduce by exploiting the honey bee life cycle. Honey bee larvae develop inside the brood cells in the wax comb. When the larvae are ready to transform from larvae to adults (metamorphosis), adult worker bees will seal larvae (now pupae) inside individual brood cells. The larvae will pupate and emerge as adults.

Female Varroa mites enter the bee hive's brood cells to reproduce. The mites use the same chemical cues that bee larvae produce that signal worker bees to seal the cell (Figure 2). The Varroa female punctures the underside of the abdomen of the pupa to feed on the fat body cells before laying its first egg.

Like honey bees, the male mite develops from an unfertilized egg. After about 72 hours in the cell, the female mite lays an egg that becomes a haploid male. A haploid individual has only one set of parental chromosomes. The female mite continues to lay about one egg per day. These eggs develop into daughters that must mature and mate with the male before the bee emerges from the brood cell. Only mature female mites will survive after leaving the brood cells. The immature mites and males die soon after the adult bee emerges.




Figure 2. **A.** An adult female mite in a honey bee brood cell before it is sealed. **B.** The pupa has been removed from this brood cell to reveal mites and a white patch of mite feces. Mites gather to mate in the fecal patch.

Varroa mites prefer to enter drone cells but can also reproduce in the worker cells of European honey bees. In Asian honey bee colonies, the mites are only able to reproduce in drone cells. Varroa mites are now so common that they can be found in nearly every hive in the United States.

Symptoms of Infestation

Heavily infested bee hives can look very healthy and produce good honey crops, only to dwindle and die during fall or winter. Colonies that have large bee populations produce a lot of brood, which allows mite populations to grow.

The Varroa mite population in the hive usually peaks in early fall. This is when the bees are getting ready to produce the last batch of worker bees for the year. These “winter bees” do not need to make protein-rich royal jelly to feed larvae so the pollen that the young bees eat is converted into vitellogenin, the protein that is stored in their fat bodies and blood (hemolymph), which helps them live all winter long. Winter bees also use vitellogenin to make royal jelly that they feed to larvae as the queen starts laying eggs even before the flowers resume blooming in the spring.

Winter bees also need to produce heat to incubate the brood at about 93°F when egg laying resumes (usually about

February). It is critical for the hive to have a healthy winter bee population for colony survival. Varroa mite feeding causes vitellogenin levels to drop. The feeding mites can also transmit viral diseases, which drastically reduce the bees’ lifespans.

The symptoms of “parasitic mite syndrome” are common in hives in the fall, but can also occur earlier in the growing season or during winter (Figure 3).

Typical symptoms include evidence of a number of virus and brood diseases, but sometimes the colony just dwindles



Figure 3. This photo shows symptoms of parasitic mite syndrome. Note the two worker bees with deformed wings and a mite on the bee in the center. The brood pattern is spotty because bees have removed dead larvae.

with no obvious symptoms as the sick foraging bees simply do not return to the hive. A dwindling adult population is also symptomatic of high tracheal mite infestations.

Varroa mites transmit many bee diseases. When Varroa infestations occur, beekeepers may observe brood disease, which causes larvae to die in their cells before the cells are capped. Beekeepers may also observe sacbrood, which is a virus disease that causes larvae to die about the time that the brood cell is sealed.

By far the most common disease associated with Varroa infestation is deformed wing virus (DWV). Honey bees often have DWV infections without visible symptoms. But when Varroa are present, virus concentrations rise and some bees may have deformed wings as a result of the DWV virus. The virus can grow in both the bee and the mite, and the mite transmits the virus to the bee during feeding — Bee-to-bee transmission can also occur. DWV and acute bee paralysis virus (ABPV) are often associated with mite-induced colony collapse.

Monitoring

It is important for beekeepers to have an idea of the mite populations in their hives. It is easy to overlook the presence of Varroa mites during a casual hive inspection. Beekeepers usually do not notice the mites on bees until there are a lot of mites in the hive.

Since Varroa mites prefer drone brood, mites may be found by opening sealed drone cells. Bees often build drone comb between the hive boxes, so splitting the boxes apart may open drone cells for inspection. Look for the coppery

brown mites on the white pupae. If many cells look infested, steps should be taken to control the mites.

The most sensitive method for detecting mites is to sample them with a “sticky board.” Some hive boxes have screened bottom boards and a plastic sheet can be inserted after spraying it with vegetable oil. Or, you can construct a sticky board of three-eighths-inch-thick wooden frame with a screen stapled to it. Put contact paper on the bottom, spray vegetable oil on it, and insert it in the colony entrance (Figure 4). The screen prevents the bees from removing mites from the sheet.



Figure 4. This beekeeper is inserting a homemade sticky board sampling device into a hive to monitor varroa mite populations.

It is also possible to remove mites from adult bees for sampling. The “ether roll” method consists of scooping up half a cup of bees (about 300 workers — be careful not to collect the queen) into a jar with a screened lid. Spray engine starter fluid into the jar, which will kill the bees and dislodge the mites. The mites stick to the glass.

Another popular method to check for mites is to again use a jar with a screened lid, then put half a cup of bees (about 300 bees) in the jar. Instead of killing the bees, coat them liberally with powdered sugar. Let the jar sit for two minutes and then shake the mites onto a white surface. Keep shaking until few mites are falling. Spraying water on to the powdered sugar makes the mites easier to see.

It is best to take three samples for a more accurate measure of the mite population on the bees.

When the mite population reaches a particular level, apply a treatment to kill the mites. Treatment thresholds vary with the season.

For example, if the ether roll or powdered sugar tests reveal 15 mites on about 300 bees, the hive has a 5 percent infestation rate. A spring infestation of 3 percent or a late-season infestation of 5-10 percent is unacceptably high in the Midwest. For the sticky board test, count the mites that fell on the board for a 24-hour period. Treat for mites if 10 mites are found on the board early in the spring, or 50 mites on the board late in the season.

Control

Table 1 shows most of the products that are registered

to control Varroa as of January 2017. Sucrose octanoate is also registered, but not recommended.

These control agents can be divided into three categories:

1. Synthetic miticides (such as fluvalinate, coumaphos and amitraz)
2. Naturally occurring molecules (such as formic and oxalic acids)
3. Products derived from plant volatiles (such as thymol, hop beta acids, and other plant oils).

Registration Number	Product Name	Active Ingredient
2724-406	Zoecon RF-318 Apistan Strip	fluvalinate (10.25%)
61671-3	For-Mite (fumigant)	formic acid (65.9%)
73291-1	Api Life Var (fumigant)	thymol (74.09%), oil of eucalyptus (16%), menthol (3.73%)
75710-2	Mite-Away Quick Strips (fumigant)	formic acid (46.7%)
79671-1	Apiguard (fumigant)	thymol (25%)
83623-2	Hopguard II (fumigant)	hop beta acids resin (16%)
87243-1	Apivar (strip)	amitraz (3.33%)
91266-1	Oxalic Acid Dihydrate (fumigant or liquid dribble onto bees)	oxalic acid (100%)
11556-138	Checkmite+Bee Hive Pest Control Strip	coumaphos (10%)

Beekeepers administer synthetic miticides by hanging a miticide strip between the combs. For the other treatments (except dribbling oxalic acid onto clusters of bees), beekeepers administer them as vapors (or fumigants) within the hive. It is important to follow product directions and pay careful attention to temperature and hive ventilation.

Many U.S. mite populations have become resistant to fluvalinate and coumaphos, but these products may regain their effectiveness if they have not been used for several years. Amitraz is a good choice for commercial beekeepers who have large numbers of hives and need a reliable and easy-to-apply control method. The other products listed will provide adequate control when used properly, but may require repeated applications especially when brood is present.

When they can access brood cells, most of the Varroa mites in a hive will be sealed inside the brood cells, so the miticide will not kill them. Worker brood is sealed for about two weeks before the adult bees emerge. Be sure to wear protective

gloves and follow label instructions. To avoid contaminating honey, some of these products should **not** be applied when honey supers are on the hive.

Mite-tolerant Honey Bees

Honey bees are gaining some ability to fight back against Varroa mites. The two most important traits that allow bees to control mites are both behavioral responses. The first trait involves uncapping and removing mite-infested pupae. Bees with this trait have what is called Varroa-sensitive hygiene (VSH). Researchers believe VSH behavior is probably triggered by the odors that are given off by the pupae that have been fed upon by mites and are infected with high levels of virus.

Some mites are infertile and these mites do not feed on pupae. Bees seldom open brood cells with mites that do not lay eggs. The USDA Baton Rouge Bee Lab has selected for this trait and made VSH bees available to queen producers. The Baton Rouge Bee Lab has also imported bees from Russia that have developed some ability to limit mite populations. It is not clear how the Russian bees do this but it is at least partly a result of the bees' behavior.

"Selecting for Varroa Sensitive Hygiene" from eXtension describes methods for assessing VSH behavior in honey bees, <<https://articles.extension.org/pages/30984/selecting-for-Varroa-sensitive-hygiene>>.

The other trait that helps bees get rid of mites is grooming behavior. Some bee stocks are better at removing mites from their bodies than others. This behavior is also linked to mite-biting. Colonies that groom more mites from themselves tend to have a higher proportion of chewed mites on the sticky board sampling sheets.

Grooming behavior may interfere with monitoring mite levels with a sticky board because the bees are actively removing mites from themselves, but in general, most mites fall passively as adult bees emerge from brood cells or bees rub against each other in the hive, which dislodges mites. You can determine if mites from sticky boards have been chewed by putting them on their backs and examining them at 15X magnification. This allows you to see the legs, which are more susceptible to chewing damage. A high proportion of chewed mites indicates relatively high grooming activity in the hive.

The Purdue University bee lab has conducted a breeding program to increase the level of mite grooming in bees, and some queens from these stocks are available from queen producers in the Midwest. Other sources of mite-grooming bees are being identified as queen producers become aware of the value of this trait.

When deciding where to purchase new queens, try to find sources that advertise mite tolerance or are produced in your region. Using local queens usually means that you must wait until June to purchase bees. However, remember that even mite-tolerant bee stocks can eventually collapse from

heavy Varroa mite infestations, so it is important to continue monitoring mite populations and use miticide when needed. Remember that when more susceptible colonies collapse from mite infestations, the mites from those colonies will end up in other nearby hives. By keeping an eye on mite levels and using mite-tolerant stock, it is possible to keep winter hive losses to about 15 percent or less.

TROPILAEELAPS MITES: A TROPICAL PEST OF HONEY BEES

Two of the four known species of *Tropilaelaps* mites (*T. clareae* and *T. mercedesae*) have made the host shift from the giant honey bee in Asia (*A. dorsata*), to our western honey bees, *A. mellifera*. *Tropilaelaps* mites are mostly restricted to the tropics in Asia and in one location in Africa. The Animal Plant Health Inspection Service (APHIS) is currently monitoring each year for *Tropilaelaps* mites in the U.S. because they can be devastating to western honey bees in the tropics. As of this writing, these mites have not been found in the U.S., Australia, South America or Europe.

Tropilaelaps feed and reproduce only on bee larvae in brood cells. They damage bees in a way that is similar to Varroa because they transmit viruses during feeding, such as deformed wing virus. *Tropilaelaps* are much smaller and can move much faster than Varroa (Figure 5). Their mouth parts are not large enough to allow them to feed on adult bees and these mites usually die within 3 days when no brood is present to feed on or to lay eggs upon. A break in the brood cycle occurs in temperate climates as the queen honey bee stops egg laying as winter approaches. Putting bees in a screen package without brood for 48 hours has been considered sufficient to rid the bees of *Tropilaelaps*. For these reasons *Tropilaelaps* is unlikely to become a major problem in the U.S., at least outside of Florida, where honey bees can have brood year-round. But *Tropilaelaps* mites have been found in South Korea and northern China, which have temperate climates. At this time, there is no reason for beekeepers in the Midwest to be concerned about this pest.



Figure 5. A Varroa mite on the left and a *Tropilaelaps* mite on the right. (Photo Credit: Zachary Huang)

READ AND FOLLOW ALL LABEL INSTRUCTIONS. THIS INCLUDES DIRECTIONS FOR USE, PRECAUTIONARY STATEMENTS (HAZARDS TO HUMANS, DOMESTIC ANIMALS, AND ENDANGERED SPECIES), ENVIRONMENTAL HAZARDS, RATES OF APPLICATION, NUMBER OF APPLICATIONS, REENTRY INTERVALS, HARVEST RESTRICTIONS, STORAGE AND DISPOSAL, AND ANY SPECIFIC WARNINGS AND/OR PRECAUTIONS FOR SAFE HANDLING OF THE PESTICIDE.

September 2018

It is the policy of the Purdue University Cooperative Extension Service that all persons have equal opportunity and access to its educational programs, services, activities, and facilities without regard to race, religion, color, sex, age, national origin or ancestry, marital status, parental status, sexual orientation, disability or status as a veteran. Purdue University is an Affirmative Action institution. This material may be available in alternative formats.

This work is supported in part by Extension Implementation Grant 2017-70006-27140/ IND011460G4-1013877 from the USDA National Institute of Food and Agriculture.